

Self-Replicating Functions

Tyler Neylon

620.2017

These are notes I'm creating for myself as I explore functions f that can be written as a sum $f = g_1 + g_2$ where g_1 and g_2 are shifted and possibly reflected versions of each other, both strongly resembling the original function f . When a function f has these properties, I informally call it a *self-replicating function*.

Like the word *fractal*, this term is not rigorously defined — in particular, it depends on the ambiguous notion of “strong




Figure 1: As an example of a self-replicating function, the normal curve can be expressed as the sum of two normal-like curves that are reflections of each other.

resemblance” — although I plan to investigate more precise requirements below.

[These notes are available in several formats: [html](#) | [standard pdf](#) | [kindle-friendly pdf](#). The source files, in LaTeX/markdown format, are available [on github](#).]

1 Motivation

I became interested in self-replicating functions by working on algorithms to procedurally generate 3d models of natural-looking trees. When algorithmically making trees, it makes sense to start from the idea of an *L-system*, which can be visualized as a kind of fractal in which a trunk forks into branches that themselves fork into smaller subbranches, this process repeating infinitely.

I noticed that tree-like *L-systems* can have a large amount of “branch overlap” concentrated around a central area of their apparent surface. For example, consider the two images in figure 2. On the left is a standard *L-system* along with a histogram showing the density of leaf points along the edge. Intuitively, the leaf points are dense even toward the extreme angles of the tree’s top. However, the density increases continuously toward the center.

We could think of each leaf point as doing a certain amount of work by covering some area along the top of the *L-system*. Each subtree is so oblivious to its other subtrees that they overlap heavily, and the central leaf points end up being highly redundant.




Figure 2: Left: An L -system; Right: the same system with two large subtrees removed. In both cases, a histogram of leaf point density is provided around an outer ellipse.

To illustrate this redundancy, the right-hand figure shows the exact same L -system with essentially half of the tree removed — yet the shape formed by the leaf points is only slightly changed.

One approach to smoothing out the distribution of leaf points would be to compromise the fractal-like nature of the system by choosing each line direction based on where it is within the fractal, rather than simply by making each branching point a smaller version of its parent. The line directions can be chosen so that the set of points at a fixed distance from the trunk point form a set of equidistant angles from a central point. The result is an extremely regular edge, as seen in figure 3.

This is ideally efficient in that each leaf point is equally important in forming the shape of the system. However, this system is defined in terms of the path to each point. Is it possible to




Figure 3: A L -like system in which line directions are chosen to maximize the regularity of leaf point distribution.

design a system so that the overall distribution of leaf points is fairly even, yet each subtree's shape is independent of its position within the full tree?

If this goal were achieved, we would necessarily have a leaf point distribution which was the sum of two smaller versions of itself. Intuitively, the leaf-point distribution of any L -system is already a self-replication function because, if its two main subtrees have distribution functions g_1 and g_2 , then the full tree has distribution function $f = g_1 + g_2$. I have to say *intuitively* here because I haven't formally defined the leaf-point distribution of an L -system.

Thus, L -systems naturally coincide with self-replicating functions. Although there are probably self-replicating functions which do

not correspond with L -systems, I nonetheless find it interesting to independently explore the world of self-replicating functions.

2 Simple cases

Technically, any polynomial can be seen as a kind of self-replicating function. For example, if $f(x) = 2x^2$,

$$\begin{aligned}g_1(x) &= (x + 1)^2 - 1 = x^2 + 2x, & \text{and} \\g_2(x) &= (x - 1)^2 - 1 = x^2 - 2x,\end{aligned}$$

then $f = g_1 + g_2$, and each g_i is a scaled shift of the original function f . In general, if $f(x) = ax^n + O(x^{n-1})$ then we can choose $g_i(x) = a/2(x \pm 1)^n + O(x^{n-1})$ so that $f = g_1 + g_2$, and each g_i has

$$\lim_{x \rightarrow \pm\infty} \frac{2g_i(x)}{f(x)} = 1,$$

which is good enough for me to subjectively say that they strongly resemble shifts of f .

However, the original motivation for self-replicating functions is based on distribution functions, so the rest of this note focuses on functions f for which $\lim_{x \rightarrow \pm\infty} f(x) = 0$.

Another simple approach would be to set $g_1 = g_2 = \frac{1}{2}f$ for any function f . This is not very interesting, and the word *shifted* in

the informal definition of a self-replicating function is intended to defeat this trivial case. That is, each g_i is expected to be similar to a translation of f , such as $f(x - 1)$ or $f(x + 1)$.

2.1 Indicator functions

The next function I'll describe is simple and meets all of the requirements so far. An *indicator function* is a function taking on only the value 0 or 1; it's also sometimes referred to as a *characteristic function*. If f is an indicator function, then you can think of those x with $f(x) = 1$ as belonging to the subset of the domain which is *indicated* by the function. It's handy to use the following bracket notation of Knuth and others: given any boolean predicate $P(x)$, let $[P(x)]$ denote the value 1 when $P(x)$ is true, and false otherwise (Knuth 1998).

Given a half-open interval $[a, b)$, define $I_{[a,b)}$ to be the function $[x \in [a, b)]$. The following equation shows how such indicator functions can be considered simple self-replicating functions:

$$I_{[0,2)} = I_{[0,1)} + I_{[1,2)}.$$

In order to match the equation $f = g_1 + g_2$, emphasizing the similarity between the g_i 's and f , we can set $f = I_{[0,2)}$, $g_1 = f(2x) = I_{[0,1)}$, and $g_2 = f(2(x - 1)) = I_{[1,2)}$.



Figure 4: Visual representation of the addition of indicator functions of intervals.

2.2 Ramp functions

Things get more interesting when $g_1(x)g_2(x) \neq 0$ for some x . To this end, define the *ramp function* for values a, b, c, d with $a < b < c < d$ via

$$J_{a,b,c,d} = \begin{cases} \frac{x-a}{b-a} & \text{if } x \in [a, b), \\ 1 & \text{if } x \in [b, c), \\ \frac{d-x}{d-c} & \text{if } x \in [c, d), \text{ and} \\ 0 & \text{otherwise.} \end{cases}$$

Then $J_{0,1,4,5} = J_{0,1,2,3} + J_{2,3,4,5}$, as illustrated in figure 5.

The ramp function example gives me four ideas for further study:

1. The addends and the sum cannot be expressed as linearly related; that is, there is no linear function $\ell(x)$ so that $J_{0,1,2,3}(\ell(x)) = J_{0,1,4,5}(x)$. Contrast this with the interval functions where $I_{[0,1)}(x/2) = I_{[0,2)}$. This raises the




Figure 5: Visual addition of two ramp functions to form another.

questions: Which self-replicating functions allow for this linear-relation restriction? Is there a slight modification of ramp functions which meets this linear-relation restriction?

[This question isn't answered in these notes.]

2. The ramp functions are piece-wise linear, but that linearity is not really the key to their being self-replicating. Rather, the key is that the left ramp and right ramp sum to 1, which matches the middle height of the functions. Which more general self-replicating functions can be constructed using this idea?

[This question is answered below.]

3. What happens if we treat the sum $f = g_1 + g_2$ as part of a sequence? Thinking of L and R for *left* and *right*, let $f_L^{(0)} = J_{0,1,2,3}$, and $f_R^{(i)} = f_L^{(i)}(x - 2)$ for $i \geq 0$. Thinking of S for *sum*, define $f_S^{(i+1)} = f_L^{(i)} + f_R^{(i)}$ for $i \geq 1$. If $f_L^{(i)}$ is positive on $(0, b)$, then $f_R^{(i)}$ is positive on $(2, b + 2)$, so $f_S^{(i+1)}$ is positive on $(0, b + 2)$. Set $f_L^{(i+1)} = f_S^{(i+1)}(x(b + 2)/b)$ so

that we maintain the region on which the left function is positive. In this way, we get a sequence of functions. What is the limiting behavior? Can we attempt to extend the sequence backwards? Can we say anything in general about the limiting behavior of a class of starting functions $f_L^{(0)}$?

[This question isn't explicitly answered below, but I suspect these notes offer some solid hints toward the solution.]

4. For the current ramp functions, the middle section is flat with value 1, while the edges sum to 1. Can we do something more interesting where the edges sum to a non-constant value? I can imagine this leading to a discontinuous function. Is there a way to do this where the functions are continuous, or at least continuous almost everywhere? Can we describe a general class of self-replicating functions which are not continuous, such as the indicator function of the Cantor dust?

[The continuous flavors of this question are answered below.]

2.3 Nonlinear ramps

Other curves that sum to 1 could easily take the place of the left and right edges of the ramp function. For example, the left and right ramps could be replaced by curves with the shapes of x^2 and $1 - x^2$ on $x \in [0, 1]$, as illustrated in figure 6.




Figure 6: An example of nonlinear ramp functions using x^2 on $x \in [0, 1]$ to determine the edge shapes.

Given any function $f : [0, 1] \rightarrow [0, 1]$, the generalized ramp function is

$$K_{a,b,c,d} = \begin{cases} f\left(\frac{x-a}{b-a}\right) & \text{if } x \in [a, b), \\ 1 & \text{if } x \in [b, c), \\ 1 - f\left(\frac{x-c}{d-c}\right) & \text{if } x \in [c, d), \\ 0 & \text{otherwise.} \end{cases}$$

If any function can be written as $K_{a,b,c,d}$ for some value of f , I'll call it a *K-function*. This form is general enough to include interval functions by using, for example, $f(x) = 0$. All previous ramp functions $J(x)$ are also *K-functions*, as can be seen by setting $f(x) = x$. Note that a *K-function* $K_{a,b,c,d}$ must have $|b - a| = |d - c|$ in order to allow a shifted version to sum to 1 throughout the middle.

The versatility of the *K-functions* shows that we can produce self-replicating functions that are highly discontinuous, such as by




Figure 7: Example K –functions: on the top is a function more discontinuous than the indicator function of an interval; on the bottom is a continuous but non-peak-monotonic function.

setting $f(x)$ to be the indicator function of a set with many border elements. Even among continuous functions, we can produce self-replicating functions which avoid being “mostly monotonic.” In particular, I’ll say that a function $f : \mathbb{R} \rightarrow \mathbb{R}$ is *peak monotonic* iff there is a point x such that $a < b < x \Rightarrow f(a) \leq f(b)$ and $x < c < d \Rightarrow f(c) \geq f(d)$. The indicator function of an interval and the ramp function are both peak monotonic, while the example K –functions in figure 7 are not.

2.4 Non-plateau functions

The ramp functions $K_{a,b,c,d}$ all have the constant value 1 on the middle interval $[b, c]$. This requires the ramps on intervals $[a, b]$ and $[c, d]$ to sum to 1. In this section, I’ll consider what can happen if we relax this condition. I’ll informally call these *non-plateau functions*.

It will be useful to propose one possible formalization of a self-replicating function before exploring non-plateau functions.

2.4.1 A formal definition for self-replicating functions

I think the term *self-replicating function* is best left as an intuitive, non-rigorous concept because there seem to be a wide variety of instances that are best studied via their own particular flavors of a formal definition. A number of other terms used to discuss mathematics are similarly unformalized or context-specific: consider *fractal*, *symmetry*, or *closure* as examples. Nonetheless, many self-replicating functions meet the conditions of the definition I'll present next.

Call a function f *exactly self-replicating* iff there exist continuous bijections s , t_1 , and t_2 such that s is not the identity function and

$$\left. \begin{aligned} f_L(x) &= f(x), \\ f_R(x) &= f(s(x)), \\ f_S(x) &= f_L(x) + f_R(x), \text{ and} \\ f_L(x) &= t_2(f_S(t_1(x))). \end{aligned} \right\} \quad (1)$$

The L , R , and S subscripts are meant to hint that these functions act as the *left* addend, *right* addend, and the *sum*; the s function suggests a *shift*, while the t_1 and t_2 functions suggest a *transformation*. The last equation in (1) captures the

similarity relationship between the addend $f = f_L$ and the sum $f_S = f_L + f_R$.

Example The ramp functions given in §2.2 and §2.3, viewed as K –functions, all adhere to the general form

$$K_{0,1,2,3} + K_{2,3,4,5} = K_{0,1,4,5}.$$

In this case, $f(x) = f_L(x) = K_{0,1,2,3}$ and $f_R(x) = K_{2,3,4,5} = f(x - 2)$. We can satisfy all of the equations of (1) by using these functions:

$$\left. \begin{aligned} t_1(x) &= \begin{cases} x & x \leq 1, \\ 3x - 2 & 1 < x < 2, \\ x + 2 & x \geq 2; \end{cases} \\ t_2(x) &= x; \text{ and} \\ s(x) &= x - 2. \end{aligned} \right\} \quad (2)$$

This is a simple yet foundational case — it may be interesting to see which other functions are exactly self-replicating with these parameters.

2.4.2 Characterizing one type of exactly self-replicating function

Note: The style of writing is about to get a tad more logically precise as some proofs start to show up.

In this section I'll give sufficient and necessary conditions for a function to be exactly self-replicating with the s , t_1 , and t_2 functions given in (2), and with $f(x) = 0$ outside of the interval $[0, 3]$. This can be considered the most general version of the category of functions we've explored so far.

For convenience, I'll introduce a notation to extract a new function with domain $[0, 1]$ from any closed domain interval of an original function f . Specifically, let $f|_{[a, b]}$ denote the function with domain $[0, 1]$ where

$$(f|_{[a, b]})(x) = f(a + (b - a)x).$$

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be any function such that $f(x) = 0$ outside of $[0, 3]$. Define the functions r_L , r_R , and g via:

$$\left. \begin{aligned} r_L &= f|_{[0, 1]}, \\ r_R &= f|_{[2, 3]}, \\ g &= r_L + r_R. \end{aligned} \right\} \quad (3)$$

Conceptually, r_L and r_R are the left and right ramp functions.

I'll show that many copies of the shape of g must dominate the landscape of f in order for it to be exactly self-replicating.

Now suppose that, in addition to having $f(x) = 0$ outside of $[0, 3]$, f is also exactly self-replicating. I'll use (1) to define functions f_L , f_R , and f_S in terms of f and the functions s , t_1 , and t_2 from (2). Notice that




Figure 8: An example showing how r_L , r_R , and g are extracted from a function f , shown on top.

$$\begin{aligned} (f_S | [2, 3]) &= (f_L + f_R | [2, 3]) \\ &= r_L + r_R = g. \end{aligned}$$

Since $f_L(x) = f_S(t_1(x))$, and t_1 maps $[1\frac{1}{3}, 1\frac{2}{3}]$ to $[2, 3]$, this means $(f = f_L | [1\frac{1}{3}, 1\frac{2}{3}]) = g$ (figure 9 below may help clarify the action of t_1). Below, I'll show how repeated application of this kind of logic determines the non-ramp values of f almost everywhere; a boolean property $P : \mathbb{R} \rightarrow \{\text{true}, \text{false}\}$ is defined to be true *almost everywhere* when the set $\{x : P(x) = \text{false}\}$ has measure zero.

At this point it will be useful to begin using base-3 notation for the intervals at hand. If s is a finite string with digits from the set $\{0, 1, 2\}$, then let $0.s\bar{*}_3$ denote the closure of the set of points whose base-3 expansion begins with $0.s$. For example, $0.11\bar{*}_3$ denotes the interval $[0.11_3, 0.12_3]$ while $0.12\bar{*}_3$ denotes the interval $[0.12_3, 0.20_3]$. I'll also use $\{2\}_2^0$ to denote a digit that may be either a 0 or a 2; for example, $0.1\{2\}_2^0 1\bar{*}_3$ denotes the union of

intervals $[0.101_3, 0.102_3]$ and $[0.121_3, 0.122_3]$.

Now, instead of writing $(f | [1\frac{1}{3}, 1\frac{2}{3}]) = g$, I can write

$$(f | 1.1\bar{*}_3) = g. \quad (4)$$

It's possible to generalize this last equation so that it defines f almost everywhere on the interval $[1, 2]$.

Recall that the notation $1.\{2\}^k 1\bar{*}_3$ indicates a union of closed intervals. In the next theorem, the notation $(f | \cup_i [a_i, b_i]) = g$ indicates that, for every i in the union, $(f | [a_i, b_i]) = g$.

Theorem 1 *Suppose that f is exactly self-replicating with functions s , t_1 , and t_2 as given in (2). Also suppose that $f(x) = 0$ outside of $[0, 3]$, and that g is defined as in (3). Then, for any $k \geq 0$,*

$$(f | 1.\{2\}^k 1\bar{*}_3) = g.$$

Proof The proof is by induction on k . Equation (4) provides the base case.

For the inductive step, suppose

$$(f = f_L | 1.\{2\}^k 1\bar{*}_3) = g.$$

Then

$$(f_R | 3 + 0.\{0\}_2^k 1_{\bar{3}}) = g,$$

so that

$$(f_S | \{1, 3\} + 0.\{0\}_2^k 1_{\bar{3}}) = g.$$

Apply t_1^{-1} to the domain set of f_S to determine the corresponding domain set of f_L :

$$(f | 1.\{0\}_2^{k+1} 1_{\bar{3}}) = g.$$

□

Define the sets G_k and G via

$$G_k = 1.\{0\}_2^k 1_{\bar{3}} \quad \text{and} \quad G = \cup_{k \geq 0} G_k. \quad (5)$$

Figure 9 illustrates G_k for low values of k .

Let's check that the sets G_k are disjoint. Suppose $x_j \in G_j$ and $x_k \in G_k$, where $j < k$, and we'll work with the standard base-3 notation in which an all-2 tail is disallowed. Either $x_k = 0.\{0\}_2^k 1 \dots$ or $x_k = 0.\{0\}_2^k 2$. There are two similar cases for x_j . In the first case, the $(j+1)^{\text{th}}$ digit of x_j is 1, which is impossible for the $(j+1)^{\text{th}}$ digit of x_k . In the second case, the $(j+1)^{\text{th}}$ digit of x_j is 2 and all subsequent digits, including the $(k+1)^{\text{th}}$, are 0;




Figure 9: The inductive process in the proof of theorem 1. The top line is the domain $[0, 5]$ of f_S ; the line below that is the domain $[0, 3]$ of f ; then the subsets of $[1, 2]$ on which f is described as the induction proceeds.

this excludes equality since the $(k + 1)^{\text{th}}$ digit x_k can't be 0. In either case, $x_j \neq x_k$, confirming that G_j and G_k are disjoint.

Using this disjointedness, we can find the total measure of their union G as follows:

$$\mu(G) = \sum_{k \geq 0} \mu(G_k) = \sum_{k \geq 0} \frac{1}{3} \left(\frac{2}{3}\right)^k = 1.$$

Since each $G_k \subset [1, 2]$, this justifies the claim that theorem 1 characterizes f almost everywhere in that interval.

Readers familiar with the *Cantor set* \mathcal{C} may notice that it's closely related to the set G . In fact, \mathcal{C} is exactly the closure of $(1, 2) - G$ shifted by a unit to reside within $[0, 1]$.

What values may f take on for the points $x \in (1, 2) - G$? The

choice is still not arbitrary as the values remain related. I'll explore this question next.

Given $x \in (1, 2)$, there is some k with $x \in G_k$ iff the base-3 expansion of x contains a 1 or if it ends with the tail $\bar{0}$. This can be expressed as:

$$\begin{aligned} \text{For } x \in (1, 2), \quad x \notin G &\Leftrightarrow \\ x \in 1.\{0\}_3^\infty - \cup_{k \geq 0} 1.\{0\}_3^k \bar{0}_3. &\quad (6) \end{aligned}$$

From here on, I'll more formally use the word *expansion* — based on the idea of the base-3 expansion of a number in $[0, 1]$ — to indicate a function $E : \mathbb{N}_{\geq 1} \rightarrow \{0, 1, 2\}$ which denotes the value $v(E) = \sum_{k \geq 1} E(k)/3^k$. If $x = v(E)$, we may write $x = 0.E_3$ and think of E as an infinite string on the alphabet $\{0, 1, 2\}$.

Suppose that $f(x) = y$ for some $x \in (1, 2)$. Let E be the expansion with $x = 1.E_3$; note that E cannot be the all-zero string $\bar{0}$ nor the all-two string $\bar{2}$ since $x \in (1, 2)$. Then $f_S(x + \{0, 2\}) = y$ and, by applying t_1 , $f(x') = y$ for both $x'_1 = (x+2)/3$ and $x'_2 = (x+4)/3$. In expansion notation, we can write these last two equations as $x'_1 = x/3 + 2/3 = 0.1E_3 + 0.2_3 = 1.0E_3$ and as $x'_2 = x/3 + 4/3 = 0.1E_3 + 1.1_3 = 1.2E_3$. We can summarize this reasoning as

$$E \neq \bar{0}, \bar{2} \quad \Rightarrow \quad f(1.\{0\}_3 E_3) = f(1.E_3). \quad (7)$$

We can expand on this idea to partition $(1, 2) - G$ into subsets on which f must have the same value. To do that, it will be

useful to define the *tail* of an expansion as a way to capture end-of-string behavior. More precisely, if E is an expansion, then define $\text{tail}(E)$ via

$$\text{tail}(E) = \{y = 0.F_3 \mid \exists j, k : E(j + m) = F(k + m) \forall m \geq 0\}.$$

Intuitively, $\text{tail}(E)$ is the set of all numbers in $[0, 1]$ with the same final sequence of base-3 digits as E , ignoring any finite prefix of either expansion. For example, $x = 0.21021\overline{011}_3$ and $y = 0.001\overline{011}_3$ have $\text{tail}(x) = \text{tail}(y)$.

The following theorem builds on equation (7).

Theorem 2 *Suppose that f is exactly self-replicating with functions s , t_1 , and t_2 as given in (2). Also suppose that G is defined as in (5). Then, for $x, y \in (1, 2) - G$,*

$$\text{tail}(x) = \text{tail}(y) \quad \Rightarrow \quad f(x) = f(y).$$

Proof Note that, by (6), x and y can be expressed in expansion notation as $x = 1.E_3$ and $y = 1.F_3$ where both E and F exclude 1 from their range, and neither has an all-0 tail.

Since $\text{tail}(x) = \text{tail}(y)$, then there exist integers j, k such that

$$E(j + m) = F(k + m) \quad \forall m \geq 0. \tag{8}$$

Let p_E be the length- j prefix of E and p_F be the length- k prefix of F , and choose the expansions E' and F' so that

$$x = 1.E_3 = 1.p_E E'_3 \quad \text{and} \quad y = 1.F_3 = 1.p_F F'_3.$$

By (8), $E' = F'$. By repeated application of (7), $f(1.p_E E'_3) = f(1.E'_3)$ and $f(1.p_F F'_3) = f(1.F'_3)$. The final result is that

$$f(x) = f(1.E'_3) = f(1.F'_3) = f(y).$$

□

It turns out that theorems 1 and 2 capture *all* of the restrictions needed for f to be exactly self-replicating. This idea is formalized by the next theorem.

Theorem 3 *Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ has the value 0 outside the domain $[0, 3]$. Also suppose we're working in the context of the functions s , t_1 , and t_2 as defined in (2), and that the function g is defined as in (3). Then f is exactly self-replicating iff*

$$\left. \begin{aligned} f(1.\{2^0\}E_3) &= f(1.E_3) \quad E \neq \bar{0}, \bar{2}; \\ f(1.1E_3) &= g(0.E_3) \quad \text{for any } E. \end{aligned} \right\} \quad (9)$$

Proof The forward direction — that (9) is a consequence of f being exactly self-replicating — has already been justified by (4) and (7). Note that these last two equations are each expanded upon in theorems 1 and 2.

To verify the other direction, it will suffice to show that, if (9) is true, then so is:

$$f(x) = f_S(t_1(x)); \quad (10)$$

this equation is enough to ensure that the definition of an exactly self-replicating function, given by (1), is satisfied.

Suppose $x = 1.CE_3 \in (1, 2)$, where $C \in \{0, 1, 2\}$ and E does not have an all-2 tail; let $x' = t_1(x)$. The argument can be split into three cases based on the value of C .

Case $C = 0$: In this case, $x' = 1.E_3 \in (1, 2)$ and $f_S(x') = f(x') = f(1.E_3)$. Apply (9) to see that $f(1.E_3) = f(1.0E_3) = f(x)$, verifying (10).

Case $C = 1$: In this case, $x' = 2.E_3 \in [2, 3]$. So $f_S(x') = f(2.E_3) + f(0.E_3) = r_R(0.E_3) + r_L(0.E_3) = g(0.E_3)$. Apply (9) and continue: $g(0.E_3) = f(1.1E_3) = f(x)$. This also verifies (10).

Case $C = 2$: This cases is similar to $C = 0$, except that $x' \in (3, 4)$. Specifically, $f_S(x') = f(x' - 2) = f(1.E_3) = f(1.2E_3) = f(x)$, again verifying (10).

In all cases, equation (10) holds, ensuring that f is indeed exactly self-replicating.

□

Call the function h *tail-consistent* on a domain set A iff $\text{tail}(x) =$

$\text{tail}(y) \Rightarrow h(x) = h(y)$ for any $x, y \in A$. There's a bijection between the exactly self-replicating functions characterized by theorem 3 and an arbitrary choice of the following three functions:

$$\begin{aligned} r_L &: [0, 1] \rightarrow \mathbb{R}, \\ r_R &: [0, 1] \rightarrow \mathbb{R}, \quad \text{and} \\ h &: (1, 2) - G \rightarrow \mathbb{R}, \text{ tail-consistent } h. \end{aligned}$$

Any choice of these three functions results in an exactly self-replicating function. Given any exactly self-replicating function f with respect to s , t_1 , and t_2 given in (2), and with $f(x) = 0$ outside $[0, 3]$, there exists a unique corresponding triple r_L , r_R , and h . Theorem 3 is the key to verifying that this correspondence between such f and triples (r_L, r_R, h) is indeed a bijection.

Below is an example image depicting the function we get by choosing $r_L(x) = \cos(2x)/2 + 1/4$ and $r_R(x) = r_L(1 - x)$. The h function has the constant value $r_L(0) + r_R(0)$.

There are many exciting directions to explore from here, but I'm going to take a break now, considering this as a nice initial step into the realm of self-replicating functions.

References

Knuth, Donald E. 1998. *Fundamental Algorithms*. Third Ed. Vol. 1. The Art of Computer Programming. Addison-Wesley.




Figure 10: An exactly self-replicating function f completely determined by $r_L(x) = \cos(2x)/2 + 1/4$, $r_R(x) = r_L(1 - x)$, and the value $f(x) = r_L(0) + r_R(0)$ for all x not determined by r_L and r_R .